

Experiment 9

Acid-Base Equilibria

CH 204

Fall 2006

Dr. Brian Anderson

Last Week

Heat in chemical reactions:

heat is a measurable quantity

produced and consumed in stoichiometric amounts

Heat Capacity:

how much heat is required to raise the temperature of something by one degree Celsius (or 1 Kelvin)

Specific Heat Capacities (J/gK):

Lead 0.128

Iron 0.449

Water 4.184

This week

Weak acid titration.

Determine K_a of acetic acid by a couple different methods.

Witness the awesome power of a buffer solution to resist changes in pH.


Non-Equilibrium Reaction

Reaction goes to completion.



Equilibrium Reaction

Products react with each other to re-form the reactants.



Equilibrium Expression

For any equilibrium reaction,



products
reactants

$$K_{eq} = \frac{[C]^c[D]^d}{[A]^a[B]^b}$$

For a weak acid dissociation,



$$K_a = \frac{[H^+][A^-]}{[HA]}$$

Two Ecksepshins

Don't include *liquids* or *solids* in equilibrium expressions.




$$K_{\text{sp}} = [\text{Pb}^{2+}][\text{SO}_4^{2-}]$$




$$K_{\text{w}} = [\text{H}^+][\text{OH}^-]$$

Strong Acids



Weak Acids

All the rest!



pH of weak acids

What is the pH of a 0.1M solution of acetic acid?



	CH_3COOH	CH_3COO^-	H^+
initial	0.1	0	0
equilibrium	$0.1 - x$	x	x

Dissociation of acetic acid

	CH_3COOH	CH_3COO^-	H^+
initial	0.1	0	0
equilibrium	$0.1 - x$	x	x

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} = \frac{[x][x]}{[0.1 - x]}$$

Assume $x \ll 0.1 \text{ M}$

$$x^2 = 0.1K_a$$

$$x^2/0.1 = K_a$$

Four-Part Lab

1) Calibrate pH meter

Make sure you're in CALIBRATION mode.

Calibrate the pH meter in the order in the lab manual: **pH 7** first, then **pH 4**, then **pH 10**.

Press ENTER or CON to confirm calibration.

Last Two Titrations of Your Life

2) Titrate 25 ml of 0.1 M acetic acid using 0.1 M NaOH

DO NOT add water! No indicator this time. Titrate in a beaker, not a flask, because you need room for the pH electrode.

Record pH after the addition of every 1.0 ml of NaOH at first, and as the pH begins to change more quickly, record smaller volume increments, down to 0.2 or 0.1 ml. Try to catch points on the vertical portion of the graph.

Switch roles with your lab partner and repeat the titration a second time.

Graph pH (y-axis) versus ml added (x-axis) in Excel.

At the Equivalence Point

All of the HA has been reacted away.

If the solution was initially 0.1M acetic acid,
it is now 0.05 M acetate

At the half-equivalence point, half of the HA
has been reacted away, and the HA and
 A^- concentrations are equal.

Half-Equivalence Point

At the half-equivalence point, $[HA] = [A^-]$.

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_a = [H^+] \times \frac{[A^-]}{[HA]}$$

$$-\log K_a = -\log [H^+]$$

so when $[A^-] = [HA]$, $pH = pK_a$

A Short-Cut to K_a

- 3) Measure the pH of 1.5 M acetic acid and two buffer solutions

Use measured $[H^+]$ and known acetic acid and acetate concentrations to calculate K_a

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

Deja Vu

	CH_3COOH	CH_3COO^-	H^+
initial	1.6	0	0
equilibrium	$1.6 - x$	x	x

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} = \frac{[x][x]}{[1.6 - x]}$$

Assume $x \ll 1.5 \text{ M}$

$$x^2 = 1.6K_a$$

$$x^2/1.6 = K_a$$

And Finally...

- 4) Add strong acid & base to buffers and to water and compare the changes in pH.

pH meters need love, too

Glass bulb is very thin

Remove carefully from storage bottle – turn the bottle, not the cap

Rinse well between samples

Dab, don't wipe

Swish samples to get better reading



All Data Goes Directly into the Lab Notebook!

I'm serious.

Your TA will dock you points if he or she sees you recording data anywhere else.

I will confiscate any loose pages of data that I see in the lab, and you will have to redo that work.

Next week

Kinetics lab

Course Evaluations

TA Evaluations

Lab check-out

**If you have missed more than one lab,
e-mail me to discuss make-up week.**

Next week's quiz

pH problems again: given $[H^+]$ calculate pH
Given pH, calculate $[H^+]$

Know how to recognize a buffer solution

Know how to make up a buffer solution

Given three variables in an equilibrium expression, calculate the fourth.