


EXPERIMENT 8
THERMOCHEMISTRY

CH 204 Spring 2007
DR. BRIAN ANDERSON

THERMOCHEMISTRY

THE STUDY OF HEAT IN CHEMICAL REACTIONS.

HEAT IS PRODUCED AND CONSUMED IN CHEMICAL REACTIONS IN STOICHIOMETRIC AMOUNTS, JUST LIKE ANY OTHER REACTANT OR PRODUCT.




THERMOCHEM IN CH204

WE LOOK AT TWO DIFFERENT THERMOCHEMICAL SITUATIONS:

CALORIMETRY
ADD SOMETHING HOT TO SOMETHING COLD
HEAT LOST BY THE HOT = HEAT GAINED BY THE COLD

Hess's LAW
FORMING CHEMICAL BONDS RELEASES ENERGY.
BREAKING CHEMICAL BONDS REQUIRES ENERGY.
ADD UP ALL THE ENERGIES TO GET
THE HEAT OF REACTION, ΔH_{RXN} .



Calorimetry

CALORIMETER – A CONTAINER THAT TRAPS HEAT

PUT A KNOWN MASS OF WATER IN THE CALORIMETER, ADD SOMETHING HOT, AND MEASURE HEAT GAIN BY THE TEMPERATURE INCREASE OF THE WATER

THE CALORIMETER ITSELF ALSO HEATS UP WHEN SOMETHING HOT IS ADDED



Accounting for Heat Energy

THE TOTAL AMOUNT OF HEAT ADDED IS EQUAL TO THE AMOUNT OF HEAT ABSORBED BY THE WATER PLUS THE AMOUNT OF HEAT ABSORBED BY THE CALORIMETER: $Q_{\text{TOTAL}} = Q_{\text{WATER}} + Q_{\text{CALORIMETER}}$

THE AMOUNT OF HEAT ABSORBED BY THE WATER IS EQUAL TO THE MASS OF THE WATER TIMES ITS SPECIFIC HEAT CAPACITY TIMES THE CHANGE IN TEMPERATURE: $Q_{\text{WATER}} = m_w \times c_w \times \Delta T_C$

THE AMOUNT OF HEAT ABSORBED BY THE CALORIMETER IS EQUAL TO ITS HEAT CAPACITY TIMES THE CHANGE IN TEMPERATURE: $Q_{\text{CAL}} = C_{\text{CAL}} \times \Delta T_C$



Specific Heat Capacity

THE AMOUNT OF HEAT IT TAKES TO RAISE 1 GRAM OF A SUBSTANCE BY 1 DEGREE C

UNITS ARE J/gK

$c_{\text{subscript}}$



TONS O' VARIABLES!

$$q_{\text{TOTAL}} = q_{\text{WATER}} + q_{\text{CALORIMETER}}$$

$$q_{\text{TOTAL}} = (m_{\text{C}} \times c_{\text{s}} \times \Delta T_{\text{C}}) + (C_{\text{CAL}} \times \Delta T_{\text{C}})$$

$$\text{HEAT ADDED} = -(m_{\text{H}} \times c_{\text{H}} \times \Delta T_{\text{H}})$$

COMBINE 'EM ALL IN ONE EQUATION
AND YOU GET...



...A VERY USEFUL EQUATION

$$q_{\text{TOTAL}} = q_{\text{WATER}} + q_{\text{CALORIMETER}}$$

$$-(m_{\text{H}} \times c_{\text{H}} \times \Delta T_{\text{H}}) = (m_{\text{W}} \times c_{\text{W}} \times \Delta T_{\text{C}}) + (C_{\text{CAL}} \times \Delta T_{\text{C}})$$

$$\Delta T = T_{\text{FINAL}} - T_{\text{INITIAL}}$$

$$\Delta T_{\text{C}} = T_{\text{M}} - T_{\text{C}}$$

$$\Delta T_{\text{H}} = T_{\text{M}} - T_{\text{H}}$$

HOW ABOUT AN EXAMPLE?



THIS WEEK IN LAB

WE WILL MEASURE THE AMOUNT OF HEAT GIVEN OFF BY
50 ml of HOT WATER, BY SOME CHUNKS OF HOT
METAL, AND BY TWO CHEMICAL REACTIONS



WE'LL DO ALL THESE REACTIONS IN A
COFFEE CUP CALORIMETER.



The basic operation of calorimetry

- START WITH A KNOWN VOLUME OF A SOLUTION IN THE CALORIMETER.
- DROP IN SOMETHING HOT, OR START A REACTION THAT GENERATES HEAT.
- CLOSE THE CALORIMETER AND MEASURE THE INCREASE IN TEMPERATURE AS HEAT IS GENERATED.
 - KEEP MEASURING THE TEMPERATURE UNTIL IT FINALLY LEVELS OUT.



Fair warning

You will be collecting lots and lots o' data points, but there are no tables in the lab manual for all this data.

All time and temperature data gets recorded directly into the lab notebook. Any loose sheets of data belong to me, and you can start over.



PART ONE:

Add HOT WATER TO cold

50 mL of cold water (5°C). Add 50 mL of hot water (75°C). Final temp should be $(75 + 5) \div 2 = 40^\circ\text{C}$

BUT THE FINAL TEMP will actually be *lower* than that because the cup itself will absorb a little bit of the heat.



HEAT CAPACITY

WE WILL USE THE DATA IN PART 1 TO CALCULATE THE *HEAT CAPACITY* OF THE CUP, IN UNITS OF J/K. THIS WILL TELL US HOW MANY JOULES OF HEAT THE CUP ABSORBS FOR EVERY K (OR DEGREE C) THE CUP HEATS UP.

$$-(m_H \times c_H \times \Delta T_H) = (m_w \times c_w \times \Delta T_C) + (C_{cal} \times \Delta T_C)$$



A WORD ON HEAT CAPACITIES

SPECIFIC HEAT CAPACITY IS AN *INTENSIVE* PROPERTY. SPECIFIC HEAT CAPACITY TELLS HOW MUCH HEAT (IN JOULES) IS REQUIRED TO RAISE THE TEMPERATURE OF *ONE GRAM* OF THE SUBSTANCE BY ONE KELVIN.

HEAT CAPACITY IS AN *EXTENSIVE* PROPERTY. IT TAKES INTO ACCOUNT HOW MUCH MASS YOU HAVE.



PART 2 UNKNOWN METAL

WE WILL DETERMINE THE IDENTITY OF AN UNKNOWN METAL BY CALCULATING ITS SPECIFIC HEAT CAPACITY.

$$-(m_M \times C_M \times \Delta T_H) = (m_C \times c_s \times \Delta T_C) + (C_{cal} \times \Delta T_C)$$



UNKNOWN SUMMARY SHEET

Specific HEAT CAPACITY of YOUR UNKNOWN METAL
AND THE IDENTITY of YOUR METAL.

OBSERVATIONS ARE VALID DATA.



PARTS 3 AND 4

THE REACTIONS OF MAGNESIUM AND MAGNESIUM OXIDE
WITH HCL. MIX THESE CONTINUOUSLY, ESPECIALLY
THE MgO.

IMPORTANT: USE 2.0M HCL TO REACT WITH
THE Mg METAL (PART 3). USE 6.0M HCL TO
REACT WITH THE MgO (PART 4).

HCL IN THE HOOD IS 6.0 M.



PARTS 3 AND 4

CALCULATE HOW MUCH HEAT IS GIVEN OFF BY THE REACTION:

$$\text{HEAT added} = (m_s \times c_w \times \Delta T_c) + (C_{cal} \times \Delta T_c)$$

DIVIDE THE HEAT ADDED BY THE MOLES OF Mg OR MgO
USED TO GET ΔH IN J/MOLE



MAKING GRAPHS IN EXCEL

DRAW LINES ON THE GRAPHS YOURSELF OR HAVE EXCEL DO IT FOR YOU.



GET IT RIGHT THE FIRST TIME

- 1) START RECORDING TEMPS *BEFORE* STARTING THE REACTION
- 2) COVER AND SWIRL IMMEDIATELY!
- 3) CONTINUE RECORDING TEMPS ON THE SAME TIMELINE THROUGHOUT THE EXPERIMENT.
- 4) KEEP TAKING TEMPERATURE READINGS UNTIL THE TEMP IS CONSTANT OR DECLINING



WORKING WITH A PARTNER

PUT YOUR PARTNER'S NAME ON EVERYTHING, BUT TURN IN YOUR OWN REPORT, WITH YOUR OWN GRAPHS AND YOUR OWN UNKNOWN SUMMARY SHEET.



SOME FATHERly Advice



START THE REPORT EARLY

DON'T WAIT UNTIL ALL THE TA OFFICE HOURS
HAVE PASSED BEFORE YOU START ON THIS.

THE CALCULATIONS ARE NOT HARD, BUT
STUDENTS HAVE MORE QUESTIONS ON THIS
LAB REPORT THAN ON ANY OTHER.



THE POST-LAB

THREE CALORIMETRY PROBLEMS
TWO HESS'S LAW PROBLEMS
HINT SHEET ON THE FREEBIES PAGE
EVERYBODY GET A 10 ON THIS