


EXPERIMENT 8
THERMOCHEMISTRY

CH 204 Spring 2009
DR. BRIAN ANDERSON




LAST WEEK

DILUTIONS - BACKWARD AND FORWARD

BEER'S LAW: $A = \epsilon cl$

DETERMINED MOLECULAR FORMULA FOR CRYSTALS



THERMOCHEMISTRY

THE STUDY OF HEAT IN CHEMICAL REACTIONS.

HEAT IS PRODUCED AND CONSUMED IN CHEMICAL REACTIONS IN STOICHIOMETRIC AMOUNTS, JUST LIKE ANY OTHER REACTANT OR PRODUCT.

Thermochem in CH204

We look at two different thermochemical situations:

Calorimetry

add something hot to something cold
heat lost by the hot = heat gained by the cold

Hess's Law

forming chemical bonds releases energy.
breaking chemical bonds requires energy.

add up all the energies to get
the heat of reaction, ΔH_{rxn} .



Calorimetry

Calorimeter — a container that traps heat

put a known mass of water in the calorimeter,
add something hot, and measure heat gain by
the temperature increase of the water

the calorimeter itself also heats up
when something hot is added



Accounting for Heat Energy

The total amount of heat added is equal to the amount of heat absorbed by the water plus the amount of heat absorbed by the calorimeter: $q_{\text{added}} = q_{\text{water}} + q_{\text{calorimeter}}$

The amount of heat absorbed by the water is equal to the mass of the water times the change in temperature times its specific heat capacity: $q_{\text{water}} = m_{\text{w}} \times c_{\text{w}} \times \Delta T_{\text{c}}$

The amount of heat absorbed by the calorimeter is equal to its heat capacity times the change in temperature: $q_{\text{cal}} = C_{\text{cal}} \times \Delta T_{\text{c}}$



Specific HEAT CAPACITY

THE AMOUNT OF HEAT IT TAKES TO RAISE 1 GRAM OF
A SUBSTANCE BY 1 DEGREE C

UNITS ARE J/gK

$c_{\text{subscript}}$



A WORD ON HEAT CAPACITIES

SPECIFIC HEAT CAPACITY IS AN *INTENSIVE*
PROPERTY. SPECIFIC HEAT CAPACITY TELLS HOW
MUCH HEAT (IN JOULES) IS REQUIRED TO RAISE THE
TEMPERATURE OF *ONE GRAM* OF THE SUBSTANCE BY
ONE KELVIN.

HEAT CAPACITY IS AN *EXTENSIVE*
PROPERTY. IT TAKES INTO ACCOUNT
HOW MUCH MASS YOU HAVE.



LOTS o' VARIABLES!

$$Q_{\text{Added}} = Q_{\text{WATER}} + Q_{\text{CALORIMETER}}$$

$$Q_{\text{Added}} = (m_C \times c_s \times \Delta T_C) + (C_{\text{CAL}} \times \Delta T_C)$$

$$\text{HEAT ADDED} = -(m_H \times c_H \times \Delta T_H)$$

COMBINE 'EM ALL IN ONE EQUATION
AND YOU GET...



...A VERY USEFUL EQUATION

$$Q_{\text{Added}} = Q_{\text{WATER}} + Q_{\text{CALORIMETER}}$$

$$-(m_H \times c_H \times \Delta T_H) = (m_W \times c_W \times \Delta T_C) + (C_{\text{CAL}} \times \Delta T_C)$$

$$\Delta T = T_{\text{FINAL}} - T_{\text{INITIAL}}$$

$$\Delta T_C = T_M - T_C \quad (\text{always positive})$$

$$\Delta T_H = T_M - T_H \quad (\text{always negative})$$



This week in lab

WE WILL MEASURE THE AMOUNT OF HEAT GIVEN OFF BY 50 ml OF HOT WATER, BY SOME CHUNKS OF HOT METAL, AND BY TWO CHEMICAL REACTIONS



WE'LL DO ALL THESE REACTIONS IN A COFFEE CUP CALORIMETER.



The basic operation of calorimetry

- > START WITH A KNOWN MASS OF A SOLUTION IN THE CALORIMETER.
- > DROP IN SOMETHING HOT, OR START A REACTION THAT GENERATES HEAT.
- > CLOSE THE CALORIMETER AND MEASURE THE INCREASE IN TEMPERATURE AS HEAT IS GENERATED.
 - > KEEP MEASURING THE TEMPERATURE UNTIL IT FINALLY LEVELS OUT.



FAIR WARNING

YOU WILL BE COLLECTING LOTS AND LOTS O' DATA POINTS, BUT THERE ARE NO TABLES IN THE LAB MANUAL FOR ALL THIS DATA.

ALL TIME AND TEMPERATURE DATA GETS RECORDED DIRECTLY INTO THE LAB NOTEBOOK. ANY LOOSE SHEETS OF DATA BELONG TO ME, AND YOU CAN START OVER.



PART ONE:

Add HOT WATER TO cold

50 mL of cold water (5°C). Add 50 mL of hot water (75°C). Final temp should be $(75 + 5) \div 2 = 40^\circ\text{C}$

BUT THE FINAL TEMP WILL ACTUALLY BE *LOWER* THAN THAT BECAUSE THE CUP ITSELF WILL ABSORB A LITTLE BIT OF THE HEAT.



HEAT CAPACITY


WE WILL USE THE DATA IN PART 1 TO CALCULATE THE *HEAT CAPACITY* OF THE CUP, IN UNITS OF J/K. THIS WILL TELL US HOW MANY JOULES OF HEAT THE CUP ABSORBS FOR EVERY K (OR DEGREE C) THE CUP HEATS UP.

$$-(m_H \times c_H \times \Delta T_H) = (m_w \times c_w \times \Delta T_C) + (C_{cal}) \times \Delta T_C$$



PART 2
UNKNOWN METAL


WE WILL DETERMINE THE IDENTITY OF AN UNKNOWN METAL BY CALCULATING ITS SPECIFIC HEAT CAPACITY.

$$-(m_M \times c_M \times \Delta T_H) = (m_C \times c_s \times \Delta T_C) + (C_{cal} \times \Delta T_C)$$


UNKNOWN SUMMARY SHEET

SPECIFIC HEAT CAPACITY OF YOUR UNKNOWN METAL AND THE IDENTITY OF YOUR METAL.

OBSERVATIONS ARE VALID DATA. WHAT DOES YOUR METAL LOOK LIKE? IS IT MAGNETIC? WHAT IS ITS DENSITY?




PARTS 3 AND 4

THE REACTIONS OF MAGNESIUM AND MAGNESIUM OXIDE WITH HCl. MIX THESE CONTINUOUSLY, ESPECIALLY THE MgO.

IMPORTANT: USE 2.0M HCl TO REACT WITH THE Mg METAL (PART 3). USE 6.0M HCl TO REACT WITH THE MgO (PART 4).

HCl IN THE HOOD IS 6.0 M.



PARTS 3 AND 4

CALCULATE HOW MUCH HEAT IS GIVEN OFF BY THE REACTION:

$$\text{HEAT ADDED} = (m_s \times c_s \times \Delta T_C) + (C_{\text{cal}} \times \Delta T_C)$$

DIVIDE THE HEAT ADDED BY THE MOLES OF Mg OR MgO
USED TO GET ΔH IN J/MOLE



MAKING GRAPHS IN EXCEL

YOU'LL HAVE A TOTAL OF 5 GRAPHS (2 FOR PART 1,
AND 1 EACH FOR PARTS 2, 3, AND 4).

YOU WILL USE THE GRAPHS TO DETERMINE ΔT_C .

YOU CAN DRAW LINES ON THE GRAPHS YOURSELF OR
HAVE EXCEL DO IT FOR YOU.



GET IT RIGHT THE FIRST TIME

- 1) START RECORDING TEMPS *BEFORE* STARTING THE REACTION
- 2) COVER AND SWIRL IMMEDIATELY!
- 3) CONTINUE RECORDING TEMPS ON THE SAME TIMELINE THROUGHOUT THE EXPERIMENT.
- 4) KEEP TAKING TEMPERATURE READINGS UNTIL THE TEMP IS CONSTANT OR DECLINING



Working with a PARTNER

PUT YOUR PARTNER'S NAME ON EVERYTHING, BUT
TURN IN YOUR OWN REPORT, WITH YOUR OWN
GRAPHS AND YOUR OWN UNKNOWN SUMMARY
SHEET.

YOU ARE WORKING WITH A PARTNER TO COLLECT THE
LAB DATA, BUT YOU SHOULD STILL DO THE
WRITE-UP BY YOURSELF.



SOME FATHERLY Advice

YOUR LAB PARTNER IS A SCHEMING BACKSTABBER.

TRUST HIM LIKE YOU WOULD A RATTLESNAKE.

TRUST HER LIKE YOU WOULD A COBRA.



START THE REPORT EARLY

DON'T WAIT UNTIL ALL THE TA OFFICE HOURS
HAVE PASSED BEFORE YOU START ON THIS.

THE CALCULATIONS ARE NOT HARD, BUT
STUDENTS HAVE MORE QUESTIONS ON THIS
LAB REPORT THAN ON ANY OTHER.



THE POST-LAB

CALORIMETRY PROBLEMS AND HESS'S LAW PROBLEMS.



NEXT WEEK'S QUIZ

- KNOW THE CALORIMETRY EQUATION
- BE ABLE TO DO HESS'S LAW PROBLEMS
- KNOW THE DEFINITIONS OF THE TERMS

WE'RE USING THIS WEEK:

HEAT CAPACITY, SPECIFIC HEAT CAPACITY,
ENTHALPY OF FORMATION,
STANDARD STATE, HESS'S LAW